
Sport and Exercise Medicine

Ten Tests and Treatments to Question in Pediatric Sport and Exercise Medicine

by
Canadian Academy of Sport and Exercise Medicine
Last updated: March 2026

1 Don't order knee radiographs to diagnose Osgood Schlatter Disease in children.

Osgood Schlatter disease (OSD) is a clinical diagnosis, based on an appropriate history and typical physical findings. Knee radiographs do not need to be performed if there is no acute concern (such as trauma, suspicion of an avulsion fracture or red flag symptoms such as night pain, joint swelling, constitutional symptoms, etc) and patients are responding to treatment.

2 Don't order ultrasound as an initial investigation for shoulder/knee injuries in children.

Bony injury is more common in children and should therefore be ruled out when assessing injuries in the pediatric patient. Conventional radiography is the primary and usually only imaging modality necessary for evaluating shoulder and knee injuries. Ultrasound should not be ordered as part of the initial diagnostic workup; however, if patients do not respond to conservative management, additional imaging may be necessary.

3 Don't order scoliosis radiographic series for back pain.

Scoliosis radiographic series consisting of full spine standing radiographs are not indicated in the evaluation of back pain unless there are clinical signs of scoliosis (asymmetry on Adams forward bend test, asymmetry of the shoulders, etc). When clinically indicated, radiographs are a good initial diagnostic tool to assess back pain but should be limited to the area of interest to limit radiation exposure.

4 Don't order thoracic spine radiographs if there is clinical concern about scoliosis.

Guidelines from the American College of Radiology (ACR) and the Society on Scoliosis Orthopedic and Rehabilitation Treatment (SOSORT) recommend full spine standing posterior-anterior (PA) and lateral radiography on initial examination of scoliosis and using lower-dose radiography techniques when available. Thoracic spine radiographs are not sufficient to evaluate scoliosis as they may not allow for adequate visualization of the curves, especially potential lumbar components. They also do not allow for assessment of the Risser index, a measure of the degree of iliac apophysis ossification and a marker for both skeletal maturity and potential curve progression.

5 Don't order oblique radiographic views for investigation of spondylolysis.

Due to their efficacy, low cost, and low radiation exposure, 2-view plain films are the best initial study. Oblique radiographs detect less than 30% of spondylolysis lesions and pose increased radiation exposure with little to no increased benefit. Advanced imaging (MRI, bone scan +SPECT) may detect stress injury not seen on radiographs.

6 Don't order a head CT scan for minor head injuries/concussion.

A significant number of children undergo CT scans for minor head injuries, which are often concussions, exposing children to the potentially harmful effects of ionizing radiation and imposing undue costs to the healthcare system. A number of clinical decision rules for use of CT for minor head injury in children, including PECARN, CATCH and CHALICE rules, have been developed in the last two decades and have shown that children with low risk for clinically important structural brain injury, as in the setting of concussion (no focal neurological deficits, no altered mental state, etc), do not require CT imaging.

7 Don't immobilize a joint with suspected amplified pain syndrome (complex regional pain syndrome).

Complex regional pain syndrome (CRPS) is a chronic severe pain condition that involves peripheral, central and autonomic nervous system and immune system mechanisms. Immobilization of the painful area can lead to prolonged symptoms and poor outcomes and should be avoided unless required for underlying pathology such as fracture management.

8 Don't order follow-up radiographs for buckle fractures of the distal radius if there are no clinical symptoms at the time of follow-up.

Most if not all buckle fractures heal without complications. Follow up radiographs are not indicated if symptoms have resolved, as this will expose the child to unnecessary radiation.

9 Don't recommend prolonged absence from school following a concussion.

Relative cognitive rest during the first couple of days following a concussion is important to allow for recovery. For students, this may require a temporary absence from school. However, extended absences from school should be avoided to prevent academic setbacks and social isolation. Students do not need to be symptom-free to return to school. As soon as they can tolerate 30 minutes of cognitive activity at home with no more than mild, brief symptom exacerbation, they can return to school. If needed, a stepwise return to learn plan with appropriate academic supports can help students transition back to the classroom following injury.

10 Don't recommend prolonged rest without active interventions if symptoms persist >10 days following concussion.

Most people, regardless of age, recover from a concussion within 2-4 weeks. However up to 30%, particularly children and adolescents, experience persisting symptoms that require further treatment/rehabilitation. Cervicovestibular rehabilitation is an intervention that can significantly reduce persistent symptoms such as headaches, dizziness and neck pain. Cervicovestibular rehabilitation is recommended for adolescents who have had these symptoms for >10 days post-injury.

How the list was created

The Canadian Academy of Sport and Exercise Medicine (CASEM) Board approved the development of pediatric-specific sport and exercise medicine (SEM) Choosing Wisely Canada's recommendations. A small working group was created to review existing Choosing Wisely Canada recommendations. The working group then created a list of suggested pediatric-specific SEM recommendations based on existing research, experience and common practice patterns. This list was then sent to the Pediatric Interest Group of CASEM, as well as a pediatric orthopedic surgeon and a pediatric MSK radiologist at McMaster University, to seek feedback on each suggested recommendation. Following revisions based on the feedback, a national survey was conducted with CASEM's membership to solicit members' feedback for each recommendation. Further revisions were made by the working group and the list was then sent to CASEM's publication committee for review and feedback. Final edits were made and approved for submission to Choosing Wisely Canada by the CASEM Board.

Sources:

- 1 Cinci E, et al. Treatment of Osgood-Schlatter Disease: review of the literature. *Musculoskelet Surg* 2017;101:195-200. [PMID: 28593576](#).
Gregory JR. [Osgood-Schlatter Disease](#). Medscape. [Internet]. 2024 [cited 2024].
Joshi A. [Osgood-Schlatter Disease imaging](#). Medscape. [Internet]. 2022 [cited 2022].
Purushottam A, et al. Osgood Schlatter syndrome. *Curr Opin Pediatr* 2007;19:44-50. [PMID: 17224661](#).
- 2 Emery K. Imaging of Sports Injuries of the Upper Extremity in Children. *Clin Sports Med* 2006;25:543–568. [PMID: 16798142](#).
Gomez J. Upper extremity injuries in youth sports. *Pediatr Clin N Am* 2002;49:593– 626. [PMID: 12119867](#).
Jacobson JA, et al. Ultrasound of the Knee: Common Pathology Excluding Extensor Mechanism. *Semin Musculoskelet Radiol* 2017;21:102–112. [PMID: 28355674](#).
Kijowski R, et al. The Role of Ultrasound in the Evaluation of Sports Medicine Injuries of the Upper Extremity. *Clin Sports Med* 2006;25:569–590. [PMID: 16798143](#).
Naraghi AM, et al. Imaging of Athletic Injuries of Knee Ligaments and Menisci:Sports Imaging Series. *Radiology* 2016;281(1):23-40. [PMID: 27643766](#).
Wolf M. Knee Pain in Children: Part I: Evaluation. *PIR* 2016;37(1):18-24. [PMID: 26729778](#).
- 3 American College of Radiology. [ACR appropriateness criteria® low back pain](#) [Internet]. 2021 [cited 2021 Oct 7].
Bhatia NN, et al. Diagnostic modalities for the evaluation of pediatric back pain: a prospective study. *J Pediatr Orthop* 2008;28(2):230–233. [PMID: 18388720](#).
Booth TN, et al. ACR Appropriateness Criteria® Back Pain—Child. *Journal of the American College of Radiology*. 2017 May 1;14(5):S13-24. [PMID: 28473069](#).
- 4 Booth TN, et al. ACR Appropriateness Criteria® Back Pain—Child. *Journal of the American College of Radiology*. 2017 May 1;14(5):S13-24. [PMID: 28473069](#).
Jones JY, et al. ACR appropriateness criteria® scoliosis-child. *Journal of the American College of Radiology*. 2019 May 1;16(5):S244-51. [PMID: 31054751](#).
- 5 Cheung KK, et al. Pars interarticularis injury in elite athletes – The role of imaging in diagnosis and management. *Eur J Radiol*. 2018;108:28-42. [PMID: 30396669](#).
Ledonio CG, et al. Current evidence regarding diagnostic imaging methods for pediatric lumbar spondylolysis: a report from the scoliosis research society evidence-based medicine committee. *Spine deformity*. 2017 Mar 1;5(2):97-101. [PMID: 28259272](#).
Miller R, et al. Imaging modalities for low back pain in children: a review of spondyloysis and undiagnosed back pain. *J Pediatr Orthop* 2013; 33:282–288. [PMID: 23482264](#).
Tofte JN, et al. Imaging pediatric spondylolysis: a systematic review. *Spine*. 2017 May 15;42(10):777-82. [PMID: 27669047](#).

- 6 Babl FE, et al. Accuracy of PECARN, CATCH, and CHALICE head injury decision rules in children: a prospective cohort study. *Lancet* 2017;389:2393-2402. [PMID: 28410792](#).
- Dunning J, et al. Derivation of the children's head injury algorithm for the prediction of important clinical events decision rule for head injury in children. *Arch Dis Child*. 2006; 91(11):885-91. [PMID: 17056862](#).
- Kuppermann N, et al. Identification of children at very low risk of clinically important brain injuries after head trauma: a prospective cohort study. *Lancet* 2009;374:1160-1170. [PMID: 19758692](#).
- Osmond M, et al. Validation and refinement of a clinical decision rule for the use of computed tomography in children with minor head injury in the emergency department. *CMAJ* 2018 July 9;190:E816-22. [PMID: 29986857](#).
- Purcell L, et al. Sport-related concussion: evaluation and management. *Paediatr Child Health* 2014;19(3): 153–158. [PMID: 24665227](#).
- 7 Baerg K, et al. Canadian surveillance study of complex regional pain syndrome in children. *Pain*. 2022 Jun 1;163(6):1060-1069. [PMID: 34799536](#).
- Sherry, DD, et al. Short- and long-term outcomes of children with complex regional pain syndrome type I treated with exercise therapy. *Clin J Pain* 1999; 15(3): 218–223. [PMID: 10524475](#).
- Weissmann R, et al. Pediatric complex regional pain syndrome: a review. *Pediatric Rheumatology*. 2016 Dec;14(1):1-0. [PMID: 27130211](#).
- 8 Ben-Yakov M, et al. Five things to know about buckle fractures of the distal radius in children. *CMAJ*. 2016 Apr 19; 188(7): 527. [PMID: 26976961](#).
- Handoll HH, et al. Interventions for treating wrist fractures in children. *Cochrane Database Syst Rev*. 2018 Dec 19;12(12):CD012470. [PMID: 30566764](#).
- Koelink E, et al. Primary Care. Physician Follow-up of Distal Radius Buckle Fractures. *Pediatrics*. 2016;137(1):e20152262. [PMID: 26729537](#).
- Riera-Alvarez L, et al. Do wrist buckle fractures in children need follow-up? Buckle fractures' follow up. *J Pediatr Orthop B*. 2019; 28: 553-554. [PMID: 31305367](#).
- 9 Patricios JS, et al. Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport-Amsterdam, Oct 2022. *Br J Sports Med*. 2023 Jun;57(11):695-711. [PMID: 37316210](#).
- Purcell LK, et al. What factors must be considered in 'return to school' following concussion and what strategies or accommodations should be followed? A systematic review. *Br J Sports Med*. 2019 Feb;53(4):250. [PMID: 29500251](#).
- Putukian M, et al. Clinical recovery from concussion-return to school and sport: a systematic review and meta-analysis. *Br J Sports Med*. 2023 Jun;57(12):798-809. [PMID: 37316183](#).

- 10 Kontos AP, et al. A Randomized Controlled Trial of Precision Vestibular Rehabilitation in Adolescents following Concussion: Preliminary Findings. *J Pediatr.* 2021 Dec;239:193-199. [PMID: 34450120](#).
- Patricios JS, et al. Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport-Amsterdam, Oct 2022. *Br J Sports Med.* 2023 Jun;57(11):695-711. [PMID: 37316210](#).
- Reneker JC, et al. Feasibility of early physical therapy for dizziness after a sports-related concussion: A randomized clinical trial. *Scand J Med Sci Sports.* 2017 Dec;27(12):2009-2018. [PMID: 28211600](#).
- Schneider KJ, et al. Targeted interventions and their effect on recovery in children, adolescents and adults who have sustained a sport-related concussion: a systematic review. *Br J Sports Med.* 2023 Jun;57(12):771-779. [PMID: 37316188](#).

About Choosing Wisely Canada

Choosing Wisely Canada is the national voice for reducing unnecessary tests and treatments in health care. One of its important functions is to help clinicians and patients engage in conversations that lead to smart and effective care choices.

Web: choosingwiselycanada.org

Email: info@choosingwiselycanada.org

Twitter: [@ChooseWiselyCA](https://twitter.com/ChooseWiselyCA)

Facebook: [/ChoosingWiselyCanada](https://www.facebook.com/ChoosingWiselyCanada)